

## NOTE

### The Permanent Rank of a Matrix

Yang Yu\*

*Department of Mathematics, Rutgers University, Piscataway, New Jersey 08854*

*Communicated by Noga Alon*

Received March 5, 1998

[View metadata, citation and similar papers at core.ac.uk](#)

#### 1. INTRODUCTION

Recall that the permanent of an  $n \times n$  matrix is defined like the determinant, but with all  $n!$  terms having positive signs (see [8]). Define the *perrank* of a matrix  $A$  to be the size of the largest square submatrix of  $A$  with nonzero permanent. Throughout this paper  $F$  denotes a field,  $p$  denotes  $\text{char}(F)$  and  $Z_p$  denotes the finite prime field of  $p$  elements. All matrices have entries in  $F$ . We assume  $p \neq 2$  throughout. Of course when  $p = 2$  permanent and determinant coincide.

**LEMMA 1.1.** *Suppose  $A$  is an  $n \times m$  matrix with  $\text{perrank}(A) = m < n$ . Let  $(A \ x)$  be the  $n \times (m+1)$  matrix formed by  $A$  following with the column vector  $x$ . Then  $\dim\{x: \text{perrank}(A \ x) = m\} \leq m$ .*

*Proof.* It is clear that  $L := \{x: \text{perrank}(A \ x) = m\}$  is a subspace. We may assume that the submatrix of  $A$  formed by its first  $m$  rows has nonzero permanent. Then since for every  $x \in L$  the perrank of  $(A \ x)$  is not full, the projection of  $L$  to the first  $m$  coordinates is one to one. Hence  $\dim(L) \leq m$ . ■

\* This is part of the author's Ph.D. thesis, written under the guidance of Professor Jeff Kahn and supported by DIMACS. DIMACS is an NSF Science and Technology Center, funded under Contract STC 91-19999. E-mail: [yangyu@math.rutgers.edu](mailto:yangyu@math.rutgers.edu).

In this lemma if we replace perrank by rank, then  $\dim(L) = m$ ; so by induction on  $m$  we have

**COROLLARY 1.2.** *Let  $m \leq n$  and  $A$  be an  $n \times m$  random matrix with each entry chosen independently and uniformly from a finite field  $F$ . Then  $\Pr(\text{perrank}(A) = m) \geq \Pr(\text{rank}(A) = m)$ .*

In particular, if  $m = n$  we have  $\Pr(\text{per}(A) = 0) \leq \Pr(\det(A) = 0)$ . As  $n \rightarrow \infty$ ,  $\lim(\Pr(\det(A) = 0))$  is well-known (see e.g. [4]). But as far as we know,  $\lim(\Pr(\text{per}(A) = 0))$  is not known.

**THEOREM 1.3.** *For any matrix  $A$ ,  $\text{perrank}(A) \geq \text{rank}(A)/2$ .*

*Proof.* We may assume  $A$  is an  $n \times n$  nonsingular matrix and let  $m = \text{perrank}(A)$ . There exists an  $n \times m$  submatrix  $B$  with full perrank. Lemma 1.1 gives  $n - m \leq \dim\{x: \text{perrank}(B \ x) = m\} \leq m$ . ■

Theorem 1.3 is tight as shown by  $\begin{pmatrix} I & I \\ I & -I \end{pmatrix}$ . This is essentially the only tight example we know at this point.

On the other hand, for  $p > 0$  perrank is also bounded above in terms of rank: it is easy to see that  $\text{perrank}(A) \leq (p - 1) \text{rank}(A)$ . The key is, if a square matrix has a column (or row) repeating  $p$  times, then its permanent is 0. This simple and known observation also gives

**LEMMA 1.4.** *Suppose  $A$  is a matrix with a column repeating  $p - 1$  times,  $p > 0$ . Then adding a multiple of this column to any other column doesn't change  $\text{perrank}(A)$ .*

This lemma suggests that we can use elementary operations to study the permanent. (This idea has also been discovered independently by Kogan and Makowsky; see Section 3.5 of [7].) The following result (true for  $p = 3$  only) is an application of this idea.

**THEOREM 1.5.** *If  $p = 3$ , then  $\text{perrank}(A) = \text{perrank}(A^{-1})$  for any nonsingular  $A$ .*

*Proof.* Let  $T_1, T_2, T_3, \dots$  be a sequence of elementary column operations that change  $A_{n \times n}$  to  $I$ . Consider the sequence of operations  $T_1, T'_1, T_2, T'_2, T_3, T'_3, \dots$  on  $\begin{pmatrix} I & A \\ I & A \end{pmatrix}$ , where  $T_i$  is applied to the left  $n$  columns, and  $T'_i$  is the corresponding operation applied to the right  $n$  columns. We do operations in pairs so that every column in each intermediate matrix is repeated.

By Lemma 1.4,  $\text{perrank}\begin{pmatrix} I & I \\ A & A \end{pmatrix} = \text{perrank}\begin{pmatrix} A^{-1} & A^{-1} \\ I & I \end{pmatrix}$ . Set  $C = \begin{pmatrix} I & I \\ A & A \end{pmatrix}$ . It suffices to show  $\text{perrank}(C) = n + \text{perrank}(A)$ .

On the one hand, if  $B$  is an  $r \times r$  submatrix of  $A$  with nonzero permanent, then the submatrix of  $C$  generated (in the obvious sense) by the left upper  $I$  and the right lower  $B$  has nonzero permanent  $2^r \text{per}(B)$ . So  $\text{perrank}(C) \geq n + \text{perrank}(A)$ .

On the other hand let  $D$  be a square submatrix of  $C$  with nonzero permanent. Suppose  $D$  has  $r+s$  rows from  $(I \ I)$  with  $r$  of them having two ones,  $s$  of them having a single one; and  $D$  has  $t$  rows from  $(A \ A)$ . The general form of  $D$  is:

$$D = \begin{pmatrix} I_r & I_r & 0 & 0 \\ 0 & 0 & I_s & 0 \\ U_{t \times r} & U_{t \times r} & V_{t \times s} & W_{t \times (t-r)} \end{pmatrix}$$

Here  $U, V$  are submatrices of  $A$  and  $W$  is a submatrix of  $(A \ A)$ . Write  $W = (W_1 \ W_2)$  where  $W_1, W_2$  are submatrices of  $A$  and  $W_1$  has at least  $(t-r)/2$  columns. It is clear that  $(U \ V \ W_1)$  is a submatrix of  $A$ , so in particular  $n \geq r + s + (t-r)/2$ , or equivalently  $(t+r)/2 \geq r + s + t - n$ . Since  $\text{per}(U \ W) = 2^{-r} \text{per}(D) \neq 0$ , we have  $\text{perrank}(U \ W_1) \geq (t+r)/2 \geq r + s + t - n$ . So  $\text{perrank}(C) \leq n + \text{perrank}(A)$ . ■

## 2. THE ALON-JAEGER-TARSI CONJECTURE

The Alon-Jaeger-Tarsi Conjecture, first proposed in 1981, is

**CONJECTURE 2.1.** *For any field  $F$  with  $|F| \geq 4$  and any nonsingular matrix  $A$  over  $F$ , there is a vector  $x$  such that both  $x$  and  $Ax$  have only nonzero entries.*

This was proved by Alon-Tarsi [3] for every non-prime finite field, but the case  $F = \mathbb{Z}_p$  ( $p \geq 5$ ) remains open. (The case  $|F| = \infty$  is trivial.) See [9] for a matroid approach. Motivated by Conjecture 2.1, Jeff Kahn (personal communication) proposed:

**CONJECTURE 2.1.** *For any  $n \times n$  nonsingular matrix  $A$  over any field,  $\text{perrank}(A \ A) = n$ .*

Conjecture 2.2 implies Conjecture 2.1 via the polynomial argument of [3]. It also immediately implies Theorem 1.3. Actually Theorem 1.3 was motivated by Conjecture 2.2. Conjecture 2.2 is true when  $p = 3$  (see [3]) and we also verified it for  $n \leq 4$ .

**THEOREM 2.3.** *For any  $n \times n$  nonsingular matrices  $A_1, A_2, \dots, A_l$ ,*

$$\text{perrank}(A_1 \ A_2 \ \cdots \ A_l) \geq (1 - 2^{-l})n$$

*Proof.* Theorem 1.3 gives  $\text{perrank}(A_1) \geq n/2$  and Lemma 1.1 implies  $\text{perrank}(A_1 \ A_2 \ \cdots \ A_{k+1}) \geq (\text{perrank}(A_1 \ A_2 \ \cdots \ A_k) + n)/2$ . The theorem follows by induction on  $l$ . ■

This theorem implies a small result on Conjecture 2.1. Choose  $l = p - 2$  and  $A_1 = A_2 = \cdots = A_l = A$ . Then the argument of [3] gives

**PROPOSITION 2.4.** *For any  $n \times n$  nonsingular matrix  $A$  over  $Z_p$ , there is a vector  $x$  without zero entries such that  $Ax$  has at most  $n/2^{p-2}$  zero entries. In particular, Conjecture 2.1 is true if  $n < 2^{p-2}$ .*

We don't know examples of equality in Theorem 2.3 if  $l \geq 2$  and we believe the theorem is not tight. In fact, we don't know any example with  $\text{perrank}(A_1 \ A_2) < n$ . Motivated by the additive basis problem, the authors of [2] conjectured that  $\text{perrank}(A \ B) = n$  for any nonsingular  $n \times n$  matrices  $A, B$  over  $Z_3$ . (Remarks: this statement can be shown by Lemma 1.4 to be equivalent to Conjecture 4.3 (in the case  $p = 3$ ) of [2]. The additive basis problem was motivated by the nowhere-zero flow problem, see Section 3.3 of [5] for more details.)

Noga Alon (personal communication) has shown that for  $A$  chosen uniformly from the nonsingular  $n \times n$  matrices over  $F$ , Conjecture 2.1 is true almost surely as  $n \rightarrow \infty$ . We will show that Conjecture 2.2 is true almost surely in the same sense.

**LEMMA 2.5.** *For an  $m \times n$  matrix  $A$  with  $\text{perrank}(A \ A) = m < 2n$ ,*

$$\dim \left( L := \left\{ x: \text{perrank} \begin{pmatrix} x & x \\ A & A \end{pmatrix} = m \right\} \right) \leq \frac{m}{2}$$

(where of course  $x$  runs over row vectors of length  $n$ ).

*Proof.* We regard the columns of  $A$  as distinct even though  $A$  may have identical columns, so that we can label the column set of  $A$  by  $[n]$ . Then the column set of  $(A \ A)$  is naturally the multiset  $W := [n] \cup [n]$ . For any  $U \subseteq W$ , define  $|U|$  to be the cardinality of  $U$ ,  $\|U\|$  to be the number of repeated elements in  $U$  and  $A(U)$  to be the  $m \times |U|$  submatrix of  $(A \ A)$  consisting of the columns in  $U$ . For example,  $|W| = 2n$ ,  $\|W\| = n$ ,  $A([n]) = A$ .

Choose  $V \in \{U: U \subseteq W, |U| = m, \text{per}(A(U)) \neq 0\}$  such that  $\|V\|$  is maximum. We may assume that  $V = [r] \cup [s]$  (multiset union), where  $r = \|V\|$ ,  $0 \leq r \leq s$ ,  $r + s = m$ .

It suffices to show that the projection of  $L$  to the first  $r$  coordinates is one to one, since  $r \leq m/2$ . So assume  $x \in L$  and  $x_1 = x_2 = \cdots = x_r = 0$ . It is in fact enough to show that  $x_{r+1} = x_{r+2} = \cdots = x_s = 0$ , since  $x = 0$  is then immediate (This is as in the proof of Lemma 1.1).

Assume  $r < s$  (otherwise we are done). Let  $V' = V \cup \{r+1\}$ . Since

$$\|V' \setminus \{r+2\}\| = \|V' \setminus \{r+3\}\| = \dots = \|V' \setminus \{s\}\| = r+1 > \|V\|,$$

we have  $\text{per}(A(V' \setminus \{r+2\})) = \dots = \text{per}(A(V' \setminus \{s\})) = 0$  by our choice of  $V$ .

Since  $x \in L$ , the  $(m+1) \times (m+1)$  submatrix of  $\begin{pmatrix} x & x \\ A & A \end{pmatrix}$  consisting of the columns in  $V'$  has zero permanent. This means exactly that  $x_{r+1} = 0$  (using  $p \neq 2$ ). Similarly  $x_{r+2} = \dots = x_s = 0$ . ■

The next corollary gives Conjecture 2.2 almost surely for a uniformly chosen random nonsingular matrix, since  $\lim(\Pr(\det(A) \neq 0)) > 0$  (again see [4]).

**COROLLARY 2.6.** *Suppose  $A$  is an  $n \times n$  random matrix with each entry chosen independently and uniformly from a finite field  $F$ , then  $\Pr(\text{perrank}(A \ A) = n) \rightarrow 1$  as  $n \rightarrow \infty$ .*

*Proof.* Let  $A_i$  be an  $i \times n$  random matrix and  $q = |F|$ . It is clear  $\Pr(\text{perrank}(A_1 \ A_1) = 1) = (1 - q^{-n})$ . By Lemma 2.5 we have  $\Pr(\text{perrank}(A_2 \ A_2) = 2) \geq (1 - q^{-n})^2$ . By induction (and again Lemma 2.5) we have for  $i \leq n$ ,

$$\Pr(\text{perrank}(A_{2i} \ A_{2i}) = 2i) \geq (1 - q^{-n})^2 (1 - q^{-n+1})^2 \dots (1 - q^{-n+i-1})^2$$

Applying this with  $i = \lfloor n/2 \rfloor + 1$  yields the corollary. ■

We mention that Lemma 2.5 can be generalized. Suppose  $(A \ A \ \dots \ A)$ , with  $A$  repeated  $k$  times, has full perrank  $m < kn$ , and  $p > k$  or  $p = 0$ , then with  $L$  defined in analogy with Lemma 2.5,  $\dim(L) \leq m/k$ . We can use this and Lemma 1.4 to give an alternative proof of (essentially) the bound in Lemma 3.6 of [2].

Finally we have a strange result for fields of characteristic 0.

**LEMMA 2.7.** *Suppose the  $i$ th row of  $A_{m \times n}$  has at least  $2i - 1$  nonzero elements and  $p = 0$ , then  $\text{perrank}(A) = m$ .*

*Proof.* We proceed by induction, the lemma being trivial when  $m = 1$ . We label the column set of  $A$  by  $[n]$  and let  $B$  be the submatrix of  $A$  formed by its first  $m - 1$  rows. Any  $(m - 1)$ -set  $U$  of  $[n]$  corresponds to an  $(m - 1) \times (m - 1)$  submatrix of  $B$  and we use  $\text{per}(U)$  to denote its permanent. Since dividing a column by a nonzero number or permuting columns doesn't affect the problem, we may assume that the last row of  $A$  is  $(1, 1, \dots, 1, 0, 0, \dots, 0)$  with  $k$  ones.

Now suppose  $\text{perrank}(A) < m$ . For any fixed  $(m-1)$ -set  $U \subseteq [n]$ , let  $R = U \cap [k]$ ,  $r = |R|$  and  $V = U \setminus R$ . For any  $(r+1)$ -set  $T \subseteq [k]$ , the  $m \times m$  submatrix of  $A$  corresponding to  $T \cup V$  has zero permanent, that is

$$\sum_{S \subseteq T, |S|=r} \text{per}(S \cup V) = 0$$

We have an unknown  $\text{per}(S \cup V)$  for each  $r$ -set  $S \subseteq [k]$  and we have an equation for each  $(r+1)$ -set  $T \subseteq [k]$ . The coefficient matrix of this linear system is exactly the inclusion matrix  $M(k, r, r+1)$ . Kantor [6] showed that inclusion matrix always has full rank over field of characteristic 0. Since  $k \geq 2m-1 > 2r$ , the number of equations is at least the number of unknowns, so all unknowns are zero. In particular,  $\text{per}(U) = 0$ , that is,  $\text{perrank}(B) < m-1$ , a contradiction. ■

This lemma gives the first part of the following theorem, and the second part then follows via the argument of [3] and Theorem 1.2 of [1].

**THEOREM 2.8.** *Suppose the  $i$ th row of  $A_{n \times n}$  has at least  $i$  nonzero elements and  $p = 0$ , then (1)  $\text{perrank}(A) = n$ ;*

*(2) for any  $S_1, S_2, \dots, S_n \subseteq F$  with  $|S_i| = 3$ , there are  $x_i \in S_i$  such that  $Ax$  has no zero entries.*

## ACKNOWLEDGMENT

The author thanks Noga Alon for a very helpful conversation.

## REFERENCES

1. N. Alon, Combinatorial Nullstellensatz, *Combinatorics, Probability, Comput.*, to appear.
2. N. Alon, N. Linial, and R. Meshulam, Additive bases of vector spaces over fields, *J. Comb. Theory Ser. A* **57** (1991), 203–210.
3. N. Alon and M. Tarsi, A nowhere-zero point in linear mappings, *Combinatorica* **9** (1989), 393–395.
4. L. S. Charlap, H. D. Rees, and D. P. Robbins, The asymptotic probability that a random biased matrix is invertible, *Discrete Math.* **82** (1990), 153–163.
5. F. Jaeger, N. Linial, C. Payan, and M. Tarsi, Group connectivity of graphs—a non-homogeneous analogue of nowhere-zero flow, *J. Comb. Theory Ser. B* **56** (1992), 165–182.
6. W. M. Kantor, On incidence matrices of finite projective and affine spaces, *Math. Z.* **124** (1972), 315–318.
7. G. Kogan and J. A. Makowsky, Computing permanents over fields of characteristic 3: Where and why it becomes difficult, *Computational Complexity*, to appear.
8. Henryk Minc, “Permanents,” Addison-Wesley, Reading, MA, 1978.
9. R. Baker, J. Bonin, F. Lazebnik, and E. Shustin, On the number of nowhere zero points in linear mappings, *Combinatorica* **14** (1994), 149–157.